Jednorázová elektronická cigareta ELF BAR 1500 DISPOSABLE POD MANGO



ADDICTIVENESS:

This product contains nicotine which is a highly addictive substance and, therefore, is not recommended for use by non-smokers. No other ingredient in this product, aside from nicotine, is known to be addictive.

TOXICITY:

This product contains nicotine which may produce toxic effects if consumed orally, especially in infants or children. DO NOT drink this product. If swallowed, visit your general practitioner with this manual immediately. If spilled onto skin, rinse with cold water immediately and continue for 10-15 minutes.

INSTRUCTIONS FOR USE:

1. Remove all outer product packaging;
2. Remove rubber caps from the product
3. Take 10-15 puffs from the mouthpiece of the product when you would normally have a cigarette.

ELFBAR

www.elfbar.com



(EN)WARNING: THIS PRODUCT IS NOT FOR USE BY CHILDREN AND NON—SMOKERS.

This product is not recommended for use by children and pregnant or lactating women.

Please read this manual carefully before you begin to use this product. Keep this manual as it contains important information which you may need again.

INSTRUCTIONS FOR USE AND STORAGE OF THE PRODUCT:

Always ensure that the container is fully closed and in an upright position.

Store the product at a temperature between 18-30°C and out of direct sunlight.

Do not refrigerate, freeze, heat or store the product in temperature above 30°C.

POSSIBLE CONTRAINDICATIONS:
Propylene glycol:

- May produce adverse effects in individuals being treated with disulfiram or metronidazole
- Allergic reactions may occur in individuals with a history of atopy, childhood eczema or rhinitis (hay fever)
- Lactic acidosis may occur when large quantities of propylene glycol is consumed by children, pregnant women or individuals with hepatic or renal failure

Glycerol:

- Chances of dehydration are increased in elderly and already dehydrated individuals
- Caution should be observed in individuals with diabetes, as metabolised glycerol may cause minor hyperglycemia or glycosuria

POSSIBLE ADVERSE EFFECTS:
First-time e-cigarette users may experience irritation of the mouth, a dry throat or cough. If the symptoms persist for more than 48 hours or becomes severe, visit a general practitioner and report the incident details as per the guidance provided under 'Reporting adverse effects'. Nicotine withdrawal effects may also occur if using an inadequate strength to satisfy your cravings. These may include irritability, aggression, feeling low, anxiety, restlessness, poor concentration, increased appetite, urges to smoke, disturbed sleep or the lowering of heart rate.

REPORTING ADVERSE EFFECTS:
If you experience any side effects as a result of using this product, please notify the MHRA in relation to adverse effects relating to products purchased in the UK via the following URL-https://yellowcard.mhra.gov.uk/yellowcards/tobaccoreportmediator/. You may also register a safety concern with the relevant competent authority within your country.